

Sleep patterns, daytime predation and the evolution of diurnal sleep site selection in lorisiforms

Magdalena S Svensson^{1*}, K.A.I. Nekaris^{1,2}, Simon K Bearder¹, Caroline Bettridge³, Thomas M Butynski^{1,4}, Susan M Cheyne⁵, Nabajit Das^{1,6}, Yvonne A de Jong^{1,4}, Averee M Luhrs¹, Lydia Luncz⁸, Simon T Maddock^{9, 10}, Andrew Perkin^{1,11}, Elizabeth Pimley^{1,12}, Stephanie A Poindexter¹, Kathleen D Reinhardt¹, Denise Spaan^{1,13}, Danica J Stark^{14,15}, Carly R Starr¹⁶, Vincent Nijman^{1,2}

¹Nocturnal Primate Research Group, Oxford Brookes University, Oxford, UK

² Little Fireface Project, Rumah Hijau, Cipaganti, Garut, Indonesia

³ Manchester Metropolitan University, Manchester, UK

⁴ Eastern Africa Primate Diversity and Conservation Program, Nanyuki, Kenya

⁵ Borneo Nature Foundation, Palangka Raya, Indonesia

⁶ Primate Research Centre NE India, Guwahati, Assam, India

⁷ Department of Zoology, B.H. College, Gauhati University, Howly, Assam, India

⁸ Institute of Cognitive and Evolutionary Anthropology, University of Oxford, Oxford, UK

⁹ Faculty of Science and Engineering, University of Wolverhampton, Wolverhampton, UK

¹⁰ Department of Life Sciences, The Natural History Museum, London, UK

¹¹ Tanzania Forest Conservation Group, Dar es Salaam, Tanzania

¹² Department of Natural and Social Sciences, University of Gloucestershire, Cheltenham, UK

¹³ Instituto de Neuroetologia, Universidad Veracruzana, Xalapa, Mexico

¹⁴ Organisms and Environment Division, Cardiff School of Biosciences, Cardiff University, Cardiff, UK

¹⁵ Danau Girang Field Centre, c/o Sabah Wildlife Department, Sabah, Malaysia

¹⁶ Northern Gulf Resource Management Group, Georgetown, Queensland, Australia

Corresponding Author:

Magdalena Svensson
Oxford Brookes University
Nocturnal Primate Research Group
Oxford OX3 0BP, UK
m.svensson@brookes.ac.uk

Number of pages: 28

Abstract number of words: 250

Number of tables: 2

Number of Figures: 3

Running title: Sleep patterns in lorisiforms

Keywords: Sleep sites; Social organization; Strepsirhine

ABSTRACT

Objectives: Synthesize information on sleep patterns, sleep site use, and daytime predation at sleep sites in lorisiforms of Asia and Africa (ten genera, 36 species), and infer patterns of evolution of sleep site selection.

Materials and methods: We conducted fieldwork in twelve African and six Asian countries, collecting data on sleep sites, timing of sleep and predation during daytime. We obtained additional information from literature and through correspondence. Using a phylogenetic approach, we establish ancestral states of sleep site selection in lorisiforms and trace their evolution.

Results: The ancestral lorisiform was a fur-clinger and used dense tangles and branches/forks as sleep sites. Use of tree holes and nests as sleep sites emerged ~22 Mya (range 17-26 Mya) in Africa, and use of bamboo emerged ~11 (7-14) Mya in Asia and later in Africa. Nests are commonly used by *Galagoides*, *Paragalago*, *Galago* and *Otolemur*, tree holes by *Galago*, *Paragalago*, *Sciurocheirus* and *Perodicticus*, tangles by *Nycticebus*, *Loris*, *Galagoides*, *Galago*, *Euoticus*, *Otolemur*, *Perodicticus* and *Arctocebus*, and all but *Sciurocheirus* and *Otolemur* additionally sleep on branches/forks. Daytime predation may affect sleep site selection and sleep patterns in some species of *Nycticebus*, *Galago*, *Galagoides*, *Otolemur* and *Perodicticus*. Most lorisiforms enter their sleep sites around sunrise and leave around sunset; several are active during twilight or, briefly, during daytime.

Conclusion: Variations in sleep behavior, sleep patterns and vulnerability to daytime predation provide a window into the variation that was present in sleep in early primates. Overall, lorisiforms use the daytime for sleeping and no species can be classified as cathemeral or polycyclic.

64 To understand broader evolutionary implications of sleep among vertebrates, including sleep architecture,
65 type of sleep, intraspecific variation in sleep, sleep duration, and the ecological pressures selecting for
66 sleep and sleep site selection, a comparative approach is required (Elgar, Pagel and Harvey, 1988;
67 Lesku, Roth II, Amlaner and Lima, 2006; Rattenborg, Martinez-Gonzalez and Lesku, 2009). Sleep can
68 comprise more than 50% of a primate's activity budget (Campbell and Tobler, 1984). Sleep can occur in
69 single continuous bouts or take the form of fragmented sleep with periods of non-sleep and activity amidst
70 otherwise continuous sleep bouts. Thus, knowledge of sleep site selection and sleep patterns can provide
71 valuable insights into a species' ecology, social behavior, and habitat requirements (Anderson, 2000;
72 Mueller and Thalmann, 2000; Gursky, 2003; Grow and Gursky-Doyen, 2010). Where primates choose to
73 sleep is not only related to their body size, degree of arboreality, competition, and pressure from
74 predation and/or parasites, but also to their activity pattern (Anderson, 2000; Eberle and Kappeler, 2004;
75 Lock and Anderson, 2013; Tagg, Willie, Petre and Haggis, 2013). More than 50% of primate species are
76 nocturnal, yet comparative information on the ecology of sleep is lacking for many nocturnal taxa, vital for
77 constructing scenarios about the evolution of primate sleep (Capellini, Barton, McNamara, Preston and
78 Nunn, 2008). The use of sleep sites in primates varies substantially, ranging from the ground, rocky
79 outcrops, tree branches/forks, dense clumps of herbs and lianas, sleep platforms, tree cavities and nests
80 that are self-constructed or constructed by other species. Use of nests (either self-constructed or made in
81 tree holes or hollows) and platforms as sleep sites is common among strepsirhines and great apes, and,
82 presumably, the earliest humans (Sabater, Veá and Serrallonga, 1997; Bearder et al., 2003; Fultz, Brent,
83 Breaux and Grand, 2013; Samson and Shumaker, 2015b), but are rarely used by other haplorhines.
84 Samson and Nunn (2015) distinguished these assembled nests, on the basis that for larger primates, tree
85 hollows would not be a viable sleeping option, and suggest that ancestral Paleocene and Eocene
86 primates probably had galago-like fixed point nest use. Since most monkeys do not use nests, nest use
87 must have evolved multiple times. To be able to infer potential sleep site patterns in early primates (i.e.
88 the ones for which only morphological data are available), we also must examine how body size, forelimb
89 to hindlimb ratio, and hand dexterity combine to assist living primates in their sleep site choices (Covert,
90 2002; Gebo and Dagosto, 2004).

To examine the question further, Kappeler (1998) reviewed several explanations for the use of nests and tree cavities amongst primates, especially among lemurs. Nests may serve as concealment against predators and/or provide thermoregulatory benefits to prevent heat loss, especially for small and solitary primates (Charles-Dominique and Martin, 1972). Kappeler (1998) also posited that nests and tree cavities particularly benefit species with neonates too altricial to cling to their mother's fur by allowing them to be placed in a safe location. Through phylogenetic analyses of multiple primate taxa, he concluded that the latter hypothesis received most support for nocturnal strepsirhines. Kappeler (1998) notably lacked any *in situ* study of Asian lorises [instead citing Rasmussen (1986) and Ehrlich and MacBride, (1989)]. Regarding the paucity of field data on many primate taxa, he urged further research of wild primates to understand better the evolution of sleep site selection.

Five years after Kappeler's review, Bearder et al. (2003), focusing on the African lorisiforms, also noted the scarcity of data on sleep sites and sleep patterns even though such data are vital to understanding diversity within nocturnal primates. Most of Bearder et al.'s (2003) data were based on studies conducted in the latter part of the last century. The authors found similarities among species within the same genus, but clear differences among genera.

In the twenty-first century, substantial taxonomic changes have occurred for both the African and Asian lorisiforms. First, the dwarf galagos of the genus *Galagoides* were recognized as a polyphyletic clade (Pozzi et al., 2015), and now are comprised of *Galagoides* (western and central Africa) and *Paragalago* (eastern Africa). *Paragalago* is a sister taxon to the genus *Galago*, and *Galagoides* is a sister taxon to the clade containing *Sciurocheirus*, *Otolemur*, *Paragalago* and *Galago* (Masters et al., 2017). Second, divergence among lorisiforms is estimated to be far more ancient than previously thought; for instance *Euoticus* split from other galagos ~30 Mya and *Arctocebus* split from *Perodicticus* ~23 Mya (Pozzi et al., 2015). Third, and related to the previous two points, more species are recognized (i.e. two additional species of *Perodicticus*, four *Nycticebus*, one *Galagoides*, and one *Sciurocheirus*). Fourth, researchers studying nocturnal primates have amassed substantial new field data from countries such as Angola, Cameroon, Equatorial Guinea, The Gambia, Kenya, Malawi, South Africa, Tanzania, Cambodia, India, Indonesia, Sri Lanka and Vietnam (Nekaris, 2003a,b; Nekaris and Jayewardene 2003; Butynski and De

Jong, 2004, 2007, 2017; Pimley, Bearder and Dixson, 2005a,b; Butynski, De Jong, Perkin, Bearder and Honess, 2006; De Jong and Butynski, 2009; Svensson and Bearder, 2013; Nekaris, 2014; Kenyon et al., 2014; Bersacola, Svensson and Bearder, 2015; Engelbrecht, 2016; Génin et al., 2016; Ray, Wren and Bowers, 2016; Kumara, Sasi, Chandran and Radhakrishna, 2016; Kappeler et al., 2017). Fifth, primatologists working on diurnal primates have taken an interest in certain lorisiforms, as lorisiforms share sleep sites with diurnal primates (Llorente, Sabater and Houle, 2003), or are hunted by them (Nishida, Uehara and Nyundo, 1979; Boesch and Boesch, 1989; Pruetz and Bertolani, 2007; O'Malley, 2010; Hardus et al., 2012).

Combined, the recent advancements in our understanding of lorisiforms allow for an overview of sleep sites, sleep patterns, sleep associations, and predation pressure faced by lorisiforms while sleeping. Through the use of new genetic data on the relationships within the Lorisiformes, we predict when various sleeping patterns emerged within this group. The deep evolutionary divergence times between various lorisiform genera help us explicitly to address several questions. Do lorisiforms provide evidence that the early primate ancestors were fixed point nest users? Did nest using evolve multiple times amongst the lorisiforms? Does the ability of a neonate to cling to the mother's fur relate to the use of fixed point nests? These data can be used as a basis to understanding ancestral sleep behavior of primates that can help to inform sleep patterns that occurred later in primate evolution.

MATERIAL AND METHODS

We follow the taxonomy of Nekaris (2013a,b), but recognize the genus *Paragalago* (Masters et al., 2017), *Nycticebus kayan*, *N. bancanus* and *N. borneanus* (Munds, Nekaris and Ford, 2013), *Sciurocheirus makandensis* (Ambrose, 2013), and *Galagoides kumbirensis* (Svensson et al., 2017). We treat the Mount Kenya potto (*Perodicticus ibeanus stockleyi*) as a subspecies of *P. ibeanus*, not *P. potto* (Butynski and De Jong 2017). As such, we include 10 genera with 36 species of lorisiform. In the subsequent text, we abbreviate *Galagoides* as *Gd.* to distinguish it from *Galago* (*G.*), and *Paragalago* as *Pg.* to distinguish it from *Perodicticus* (*P.*).

144

145

Data collection

146 Post-2003 (i.e. after the publication of Bearder et al.'s 2003 compendium) we conducted nocturnal field
147 work in Angola (SKB, MSS; 1 mo), Cameroon (AML, TMB, YdJ; 3 mo), Democratic Republic of the Congo
148 (TMB; 2 mo), Equatorial Guinea (Bioko: TMB; 12 mo), Ethiopia (TMB; 1 mo), The Gambia (SKB, MSS; 1
149 mo), Kenya (TMB, YdJ; 34 mo), Nigeria (AL; 2 mo), Malawi (SKB; 1 mo), Rwanda (SKB, MSS; 1 mo),
150 Tanzania (TMB, YdJ, CB, AP; 19 mo), Uganda (TMB, YdJ, MSS, AML; 19 mo), Cambodia (CRS, KAIN;
151 11 mo), India (KAIN, ND; 32 mo), Indonesia (Java: KAIN, VN, KDR, DS; 60 mo; Sumatra: KAIN; 1 mo),
152 Malaysia (Borneo: DJS; 60 mo), Sri Lanka (KAIN, EP; 22 mo) and Vietnam (SAP, KAIN; 9 mo). We
153 collected most data on populations where individuals could not be individually recognized, but in Borneo,
154 Cambodia, India, Sri Lanka, Vietnam and Java, we followed identified individuals with radio collars or
155 other markers. We obtained additional data from published studies and through correspondence with
156 researchers, including those working on great apes (bonobos *Pan paniscus*, common chimpanzees *P.*
157 *troglydytes*, Sumatran orangutan *Pongo abelii*, Bornean orangutan *P. pygmaeus* and Tapanuli orangutan
158 *P. tapanuliensis*) to obtain data on predation events.

159

160

Analyses

161 We used species as the unit of analysis. We pooled data from studies to provide a global picture. Based
162 on previous research (Bearder et al., 2003), we placed sleep site types into five groups: nests, tree holes
163 or hollows, dense tangles of vegetation, tree branches/ forks, and bamboo thickets. We ranked the use of
164 sleep sites types from zero (no evidence of use), one (occasional use or mixed evidence) or two (regular
165 use).

166 To typify social cohesion, we collected data on sleep group size. When transporting infants, these can be
167 carried in the parent's mouth or they can cling on to their parent's fur. For each species we recorded
168 whether they carried infants in the parent's mouth or if they can cling to their parent's fur, or whether they
169 employed both methods. Regarding vocalizations, we included call types (audible to humans) used for

social cohesion, advertisement and maintenance. We exclude the ultrasonic calls of *Perodicticus*, *Loris* and *Nycticebus*. We ranked vocalizations as one (social cohesion vocalizations displayed at sleep site) or zero (social cohesion vocalizations not displayed at sleep site). Intermembral Index (IMI, a ratio of forelimb length to hindlimb length) for the different species was taken from Fleagle (2013) and for slow and slender lorises from measurements taken by KAIN and DJS on wild-caught live animals.

To gain insight into sleep patterns and the presence of fragmented sleep in the lorisiforms, we compiled data on when individuals entered and exited sleep sites. From selected sites, we added information on pre- or post-dusk waking and pre- or post-dawn sleeping. We added observations of sleep during the night or non-sleep behavior during the day.

We examined evidence of predation on lorisiforms and highlight those instances where the events occurred while the animal was asleep, or where we could reasonably infer that predation had taken place during the daytime. We excluded predation events by nocturnal predators such as owls, but included events from cathemeral or crepuscular predators. While we acknowledge that most lorisiforms, at least occasionally, sleep for brief periods during the night, and that they may be subject to predation by nocturnal predators at these times, this form of rest is distinctly different from them selecting and using a sleep site where they will sleep during day time. Additionally, we compiled information on anti-predator strategies used by lorisiforms and which of these might be most effective at sleep sites.

We carried out reconstruction of ancestral states on a subset of species for which full sleep site and fur clinging behaviour and published genetic sequences were available. We obtained cytochrome *b* sequences (1,140 bp in length) of 23 species of lorisiform from GenBank (for accession numbers see Fig. 1) and we aligned them with MAFFT v.7 multiple sequence alignment (Kato and Standley, 2013). This formed the basis of our ancestral state reconstruction analyses. We constructed phylogenetic trees using BEAST v.2.4.6 (Drummond and Rambaut, 2007; Suchard and Rambaut, 2009; Bouckaert et al., 2014). We implemented a strict clock with the birth-death speciation tree prior for 100 million generations, sampling every 10,000 iterations. We checked analyses for convergence using Tracer v.1.6. We then used the posterior probability tree produced by BEAST to perform stochastic character mapping (Huelsenbeck, Nielsen and Bollback, 2013) to infer ancestral states of sleep site and fur clinging using the

R package phytools v.0.6-20 (Revell, 2012). Phylogenetic signal was tested for discrete character evolution of each character by comparing AICc scores with and without phylogenetic error structure using the fitDiscrete function in the R package, geiger v.2.0.6 (Harmon et al., 2008). This was estimated by testing a model with complete phylogenetic independence (lambda fixed to 0) to one with phylogenetic non-independence (free lambda tree transformation).

We fitted stochastic character histories for each character set by executing continuous-time reversible Markov models, to sleep sites and fur clinging, over 99,900 simulations each. An equal rates of transition model was used to sample the state transition matrix “Q” from the posterior probability. Ancestral character estimation (“ace”) was used to demonstrate the probabilities of states at each node. To date the timing of the ancestral states of sleep site and fur clinging, the timed phylogeny of Pozzi et al. (2015) was used to calculate mean values and 95% highest probability estimates in millions of years ago (Mya).

RESULTS

Physical characteristics of sleep sites and evolution of sleep site selection

The type of sleep site lorises most commonly used was tangles (67% or 24 of 36 species), followed by holes (44%, 16 species), branches/forks (44%, 16 species), nests (either self-built or built by other species: 33%, 12 species) and bamboo (14%, 5 species) (Table 1). Of the 24 species that use tangles, 62% also use branches/forks, 46% also use holes, and 42% also use nests. Of the 16 species that use branches/forks, 94% also use tangles. Of the 16 species that use holes, 69% also use nests, 69% also use tangles, and 31% also use branches/forks. Twelve species use nests, 92% of which also use holes and 83% also use tangles. Of the five species that use bamboo, the four Asian lorises also use branches/forks and tangles, but none use nests or holes, whilst *Gd. demidovii* mainly uses nests in dense undergrowth, and, to some extent, tree holes and tangles. Species in which infants cling to the adult's fur do not tend to use nests or tree holes.

The IMI ranges from lows of around 50 in *Galago* spp, representing clear vertical clingers and leapers with legs twice as long as their arms, to an intermediate value of around 70 in *Otolemur* and *Galagoides*,

and highs of over 90 in *Loris* and *Nycticebus*, with arms and legs being almost the same length. Species with low IMIs tend to be the ones where the infants cling on the adult's fur, and that use nests and tree holes.

At least three lorisiforms use human-made sleep sites. *Galago senegalensis* sleeps in traditional bee-hives (hollowed tree boles), birdhouses, and roofs of buildings while *G. moholi* uses ventilation pipes at some study sites. *Otolemur crassicaudatus* sleeps in traditional bee-hives and roofs of buildings, and *Pg. cocos* is also known to utilize human-made sleep sites.

Phylogenetic relationships showed strong support for all splits except for the sister group relationship between *Artocebus* + *Perodictus* and *Nycticebus* + *Loris* (bpp = 0.63) (Fig. 1 and 2). Fur clinging and some sleep sites show strong phylogenetic signal under a lambda transformation model: fur clinging (estimated lambda = 1, AICc = 25.175) is a better fit (Δ AICc = 22.08) than a model with no phylogenetic signal (lambda fixed to 0, AICc = 47.255); tree hole (estimated lambda = 1, AICc = 33.64) is a better fit (Δ AICc = 19.1) than a model with no phylogenetic signal (lambda fixed to 0, AICc = 52.74); branches/forks (estimated lambda = 0.98, AICc = 28.96) is a better fit (Δ AICc = 2.948) than a model with no phylogenetic signal (lambda fixed to 0, AICc = 31.91). Nests showed some support for phylogenetic signal (estimated lambda = 0.557, AICc = 47.01) is a better fit (Δ AICc = 0.744) than a model with no phylogenetic signal (lambda fixed to 0, AICc = 47.75). Two sleep sites showed no support of phylogenetic signal: bamboo (estimated lambda = 0.363, AICc = 35.559) is a worse fit (Δ AICc = -0.144) than a model with no phylogenetic signal (lambda fixed to 0, AICc = 35.415); dense tangle (estimated lambda = 0.607, AICc = 44.78) is a worse fit (Δ AICc = -1.462) than a model with no phylogenetic signal (lambda fixed to 0, AICc = 43.318). Although stochastic character histories were estimated for all datasets, no information about ancestral evolution should be drawn from sleep sites in bamboo and dense tangles and the use of nests should be interpreted very loosely due to a lack of signal (*these were included in figures for visual representation purposes*).

At ~40 Mya (range 36-44 Mya, nb. all dates used herein are taken from Pozzi et al., 2015), the ancestral lorisiform infants were carried by clinging to the fur of its parent (Fig. 1). This ancestral state is retained in all Asian taxa as well as in some African taxa (e.g. *Perodicticus*, *Artocebus*, and *Otolemur*). Carrying

infants in the mouth evolved ~22 (17-26) Mya in the ancestor of the African galagos. The ancestral lorisiform used dense tangles and branches/forks as sleep sites. Almost all extant species still use dense tangles as sleep sites, but this trait got lost twice in the east African coastal *Paragalago* species. While the majority of species still use branches/forks as sleep sites, this trait changed at ~14 (12-18) Mya for *Paragalago* and ~12 (8-15) Mya, when *Otolemur* and *Sciurocheirus* split from the other galagos (Fig. 2).

[FIGURE 1 HERE]

The use of nests is restricted to the African lorisiforms and likely emerged ~22 (17-26) Mya, after *Euoticus* split from other galagos (Fig. 2). Use of bamboo as a sleep site appears to have emerged early on in their evolution at ~11 (7-14) Mya, after *Nycticebus* split from *Loris*. At present, all *Nycticebus* species, apart from *N. menagensis*, are known to use bamboo as sleep sites. Independently, *Gd. demidovii* uses bamboo as a sleep site but bamboos are absent over most of its geographic range; this behavior probably emerged in the last 5 million years.

[FIGURE 2 HERE]

Sleep patterns

Most lorisiforms enter their sleep site between 0.5 hr before and 0.5 hr after sunrise, and leave their sleep site between 0.5-1.0 hr before and 0.5-1.0 hr after sunset. Several lorisiforms are active (moving, feeding and calling) during twilight: e.g. *N. javanicus* and *O. garnettii* up to 1.5 hrs before sunset, and *S. alleni*, *Pg. cocos*, *Pg. zanzibaricus* and *G. senegalensis* up to 1.0 hr before sunset.

In Africa, the number of daylight hours (time between sunrise and sunset), and thus the numbers of hours available for sleep, varies between ~13 hrs (June) and ~11 hrs (December) in Senegal and Eritrea, ~10 hrs (June) and ~14 hrs (December) in southern Africa, and ~12 hrs (year round) in East Africa (Kenya, Tanzania and Uganda). We found no evidence that species in the more northern or southern regions adjust their sleep pattern. In general, for most species, sleep is an equitable 12 hrs year-round. In Asia, *N. bengalensis* in northeastern India, Myanmar and China, have ~10 hrs of daylight available for sleep in December and ~14 hrs in June; again, there is no evidence to suggest that they adjust their sleep pattern. The southern-most populations of lorisiforms in Asia are found in Sri Lanka (*Loris tardigradus*) and Java

(*N. javanicus*), both situated ~7° north and south of the equator, respectively. As such, annual variation in daylight hours is small and sleep is equitable 12 hrs year-round.

Numerous lorisiforms, including *N. javanicus*, *G. gallarum*, *G. senegalensis* and *O. garnettii* are sometimes active during the day, presumably only for short periods and possibly in response to being disturbed by humans, adverse weather or because of (real or perceived) predator threats. *Galago senegalensis* occasionally sleep in the middle of the night, but the lengths of these sleep bouts remain unknown. Additionally, *G. moholi*, *N. javanicus* and *N. pygmaeus* occasionally sleep during the night. These species have been known to use daily and multiday torpor, which may suggest they are indeed in a state of torpidity, and not sleeping (Nowack, Mzilikazi and Dausmann, 2013a; Ruf, Streicher, Stalder, Nadler and Walzer, 2015; Reinhardt, Wirdateti and Nekaris, 2016). Overall, however, the daytime is used for sleeping and we could classify no species as cathemeral or polycyclic.

[TABLE 1 HERE]

Predation at sleep sites

Predation avoidance appears to be a main factor in sleep site choice. Benefits are associated with all the sleep site types regarding protection against predation. Known predators of lorisiforms include a wide range of species, including those that target lorisiforms at their sleep sites (Table 1). Snakes and monitor lizards can access tree holes and branches/forks, whereas monkeys and apes, and possibly also some snakes, can access tree holes and tangles. Among reptiles, monitor lizards *Varanus* spp. and reticulated python *Malayopython reticulatus* prey on *N. pygmaeus* and *N. coucang*. The smoothness of bamboo stems may provide protection for *Nycticebus* spp. and *Gd. demidovii*.

Diurnal raptors prey on lorisiforms, although recorded captures are scarce. Predators known to prey on lorisiforms are: crowned eagles *Stephanoaetus coronatus* on *P. potto* and *Galago* spp., Verreaux's eagle *Aquila verreauxii* on *G. moholi*, and changeable hawk-eagle *Nisaetus cirrhatus* on *N. coucang*. These captures likely took place during the day when the lorisiforms were at their sleep site.

Small mammalian carnivores, such as palm civets, linsangs and genets (Viverridae), may capture lorisiforms when they enter or leave their sleep sites. Remains from *P. ibeanus* have been found in

leopard *Panthera pardus* scats and African palm civets *Nandina binotata* are known predators of *P. edwardsi*.

Blue monkeys *Cercopithecus mitis* prey on *Gd. thomasi* and/or *G. matschiei*, with predation observed in the afternoons. Sooty mangabeys *Cercocebus atys* have been observed poking *Gd. demidovii* out of their nests with sticks. *Nycticebus hilleri* has been observed to be captured and killed during daytime by *P. abelii* – we obtained confirmation from two sites (Table 2). Data from five sites in Borneo suggest that *P. pygmaeus* do not prey on *Nycticebus* there. *Pan troglodytes* occasionally prey on Galagidae (Fig. 3), mainly when sleeping in tree holes, sometimes using tools such as sticks. *Pan paniscus* have been observed to force *Gd. demidovii* out of tree holes by inserting fingers into the hole and then hitting the trunk (Table 2).

Humans are probably one of the main predators of lorisiforms. This relationship is especially true in Asia where *Nycticebus* and *Loris* are taken to meet the demand for the pet and traditional medicine trades, and where specialized hunters seek out sleep sites during the day.

[FIGURE 3 HERE]

[TABLE 2 HERE]

DISCUSSION

We show that lorisiforms use a wide range of sleep sites, with most taxa sleeping in dense tangles, followed by holes and on branches/forks. Fewer species use nests and bamboo. It appears that the ancestral lorisiform would have used dense tangles, and branches/forks as sleep sites. The use of tree holes and nests as sleep sites emerged ~30 (24-36) Mya in Africa, and the use of bamboo as a sleep site emerged ~31 (23-26) Mya in Asia and later in Africa. The ability of infants to cling onto their parents' fur appears to be the ancestral condition, and carrying infants in the mouth is a derived condition and emerged in the African taxa. Our data provide support for Kappeler's (1998) hypothesis that use of nests

and tree holes is linked to having altricial infants that are not able to cling to fur, thus providing them with a relatively safe location while adults forage.

Further understanding the comparative morphology of fur clingers may help us to infer nest using behavior in the fossil record. We found a strong relationship between more generalized arboreal lorisiforms with a IMI nearer to 100 in relation to fur clinging and the use of nests. Tree hole use was limited to animals with the lowest IMI that are also vertical clingers and leapers. Functionally, animals with shorter arms, and hence lower IMI, might not be able to cling as well on tangles and branches. Such morphological adaptations are further emphasized by the presence of a *retia mirabilia* (where the arteries form vascular bundles that allows blood to flow even when the animal remains still) in *Loris*, *Nycticebus* and *Perodicticus* allowing an enhanced grip (Ankel-Simons, 2000; Congdon and Ravosa, 2016). In the fossil record IMI and the ability to engage in specialized grasping may help us to interpret the sleeping patterns and sleep site selection of extinct species such as *Carpolestes simpsoni*, that resemble the more generalized arboreal lorisiforms in this study that did not use tree holes (Bloch and Boyer, 2002).

Sleep

Site selection can be seen partly as an anti-predation strategy, depending on predator species and densities (Charles-Dominique and Martin, 1972; Anderson, 2000). Factors important in reducing daytime predation appear to be: connectivity of sleep trees, use of thorny bushes, nest hole entrance size and selection of dense tangles of lianas and undergrowth and smooth-surfaced substrate such as bamboo.

The entrance size of sleep holes used by lorisiforms tends to be no larger than is necessary for the individual to enter. This minimizes the number of predator species that are able to enter or reach inside. Selection of tree holes with suitably small entrances that only enable the strepsirrhine to enter/exit is not always possible, especially when the number of tree holes in an area is limited. As a result, holes with larger entrances are sometimes used. For example, *S. cameronensis* used tree holes with entrances of 20 cm diameter larger than necessary for it to access the hole (Pimley, 2002). More studies that systematically measure tree holes used by lorisiforms are needed.

Many species reuse sleep sites in an unpredictable order. This allows them to become familiar with the sleep site and facilitate escape during predation attempts (Di Bitetti, Vidal, Baldovino and Benesovsky, 2000; Nekaris, 2003a; Qihai, Chengming, Ming and Fuwen, 2009; Svensson and Bearder, 2013). Rotation of sleep sites makes it more difficult for predators to ambush prey (Di Bitetti et al., 2000). Other species, such as *G. gallarum*, frequently sleep on branches/forks in the center of trees or bushes among a barrier of dense thorns, and use such areas on consecutive days (De Jong and Butynski, 2004a,b; Butynski and De Jong, 2013). Sleeping in tangles of dense vegetation reduces detection from predators, provides protection from the elements and facilitates rapid escape, especially for smaller lorisiforms such as *Galagoides*, *Paragalago* and *Loris* (Kappeler, 1998). Vegetation tangles and bamboo have also been hypothesized as anti-predation strategy for *Loris* and *Nycticebus* (Nekaris, 2014). The slow and slender lorises, angwantibos and pottos are non-saltatory arboreal climbers, incapable of leaping (Sellers, 1996). This locomotion demands constant connectivity to maintain substrate contact, as well as an increased number of escape routes from predators (Voskamp, Rode, Coudrat, Wilson and Nekaris, 2014).

Researchers have found animals exposed to high levels of predation to display less time spent in sleep, while those with less disturbances experience increased sleep quality (Samson and Shumaker, 2013, 2015a). This behavior is largely due to disturbances from predators during the sleeping period, as well as a need to be more alert (Zepelin, 2000; Lima, Rattenborg, Lesku and Amlaner, 2005). More field research on sleep quality is needed in primates, to determine if different sleep site types and predation pressures influence sleep patterns.

The use of different types of sleep sites within the same species of lorisiform suggests these species may be opportunistic generalists that are able to use the range of habitat features available to them or respond to varying sleep site selection pressures. Similar variability is seen in other primate species that have access to the same types of sleep sites in different environments, but do not select them based on differences in the site characteristics (Pontes and Soarse, 2005; Duarte and Young, 2011). Despite the range of sleep site types seen across the group however, sleep patterns are mostly consistent, with all species demonstrating nocturnal, not cathemeral behavior.

377 Given that most lorisiforms live in the tropics and only a few in the subtropics, with small amounts of
378 variation in day length, most species have equal amounts of time available for sleep. The exceptions are
379 *N. bengalensis*, and the southern-most populations of *G. moholi* and *O. crassicaudatus*, which have 4 hrs
380 less available for sleep in winter than in summer (for *N. bengalensis* this is the boreal winter and for *G.*
381 *moholi* and *O. crassicaudatus* it is the austral winter, when nights are significantly longer than in the
382 summer). Several species of lorisiform, mostly those studied the longest, are active for short periods
383 during the day. Conversely, these species have been recorded to sleep for up to several hours during the
384 night. This is possibly linked to low temperatures or other adverse conditions.

385 Thermoregulation may be an important factor in sleep site selection in lorisiforms, especially for species
386 that live at high elevations and/or at high or low latitudes (Ruf, et al., 2015). Tree holes provide good
387 insulation against the cold (Schmid, 1998; Daussman et al., 2004), and buffer against heat. Nests may
388 also serve thermoregulatory requirements (Radespiel, Cepok, Zietemann and Zimmermann, 1998;
389 Lutermann, Verburgt and Rendigs, 2010; Nowack et al., 2013a). *Galago moholi* return to their sleep site
390 (usually a tree hole but also nests) earlier than usual during cold nights to huddle with other individuals
391 thus serving a thermoregulatory purpose (Bearder and Martin, 1980). The requirement for
392 thermoregulation may also explain why the smaller species of lorisiform such as *Loris*, *Galago*,
393 *Galagoides*, *Paragalago* and *Sciurocheirus* (which lose heat more quickly than their larger relatives) tend
394 to sleep together, thereby sharing body heat (Nowack, Wippich, Mzilikazi and Dausmann, 2013b; cf.
395 Eppley, Watzek, Dausmann, Ganzhorn and Donati, 2017). Some of these smaller taxa, including *Pg.*
396 *zanzibaricus* and *Gd. thomasi*, bring in fresh plant material to line their sleep site (Bearder et al., 2003).

397 Peckre et al. (2016) pointed out the need for more studies on infant carrying and its relevance to primate
398 evolution, in particular regarding the evolution of an enhanced grip. Based on nearly 20 years of new field
399 data, we help to confirm the view fur clinging is an ancestral trait in lorisiforms, and that fur clinging
400 species rarely or never use tree holes (cf. Kappeler, 1998). Lorises and pottos have a shared derived trait
401 called the *retia mirabilia*, as well as a reduced second digit (Ankel-Simons, 2000). These morphological
402 traits produce an enhanced grip that has been suggested to be an anti-predation strategy (Charles-
403 Dominique, 1977; Nekaris, 2014; Oates, 1984). Where in-depth studies were conducted on apes, the

importance of sleeping posture has proven to improve sleep quality (Samson and Shumaker, 2013; Samson and Shumaker 2015a). Similarly, clinging to branches and a strong grip is also shown to be related to continuous sleep during the diurnal period as well as a decreased frequency of measurable fragmented sleep (KAIN and KDR, unpublished data). The confirmation of nest use as a derived state in strepsirhines that evolved multiple times corresponds with the deep evolutionary divergence seen among lemuriforms and lorisiforms. We provide a novel set of data that we hope will inform further studies reconstructing aspects of primate evolution.

ACKNOWLEDGEMENTS

We thank Riset and Teknologi (Ristek), Balai Konservasi Sumber Daya Alam Bandung (BKSDA Bandung), Research Center for Biology, Indonesian Institute of Sciences (LIPI) and Konservasi Sumber Daya Alam Garut (KSDA Garut), Sabah Wildlife Department, College of African Wildlife Management, Mweka, Tanzania, Institute of Primate Research (Nairobi, Kenya) and Lolldaiga Hills Research Programme (Nanyuki, Kenya).

Funding was received from: Leverhulme Trust (RPG-084) (KAIN), People's Trust for Endangered Species (KAIN), National Geographic Society (KAIN, SKB, TMB, YdJ), Disney Worldwide Conservation Fund (KAIN), Conservation International Primate Action Fund (KAIN), Cleveland Zoological Society and Cleveland Metroparks Zoo (KAIN, VN, DJS), Primate Society of Great Britain Conservation Working Party (KAIN), Amersfoort Zoo (KAIN), Primate Conservation Inc. (KAIN, TMB, YdJ, EP), Critical Ecosystem Partnership Fund (TMB, YdJ), Zoologische Gesellschaft für Arten- und Populationsschutz (KAIN), Columbus Zoo Aquarium (KAIN, DJS), Phoenix Zoo (DJS), Shaldon Wildlife Trust (KAIN), Brevard Zoo (KAIN), Henry Doorly Zoo (KAIN), Houston Zoo (DJS), International Primate Protection League (KAIN), Frederika Bremer Förbundet (MSS), Iris Stipendiet (MSS) and Oxford Brookes University Departmental Research Scheme (KAIN, VN, MSS, SKB).

Our research was approved by Oxford Brookes University, Manchester Metropolitan University and Cardiff University and complies with the American Association of Physical Anthropologists Code of Ethics.

This project was conceived by KAI Nekaris, MS Svensson and V Nijman. MS Svensson, C Bettridge, TM Butynski, YA de Jong, AM Luhrs, L Luncz, A Perkin, E Pimley and SK Bearder collected field data on the African taxa. KAI Nekaris, SM Cheyne, N Das, D Spaan, DJ Stark, KD Reinhardt, and CR Starr collected field data on the Asian taxa. V Nijman, MS Svensson, KAI Nekaris, SA Poindexter, and S Maddock analyzed the data, and MS Svensson, V Nijman, TM Butynski, YA de Jong and KAI Nekaris wrote the paper with input from the other authors. All authors approved the final manuscript.

LITERATURE CITED

- Ambrose, L. (2003). Three acoustic forms of Allen's galagos (Primates: Galagonidae) in the Central African region. *Primates*, 44(1), 25-39.
- Ambrose, L. (2013). *Sciurocheirus makandensis* sp. nov – Makandé squirrel galago. In T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 421–422). London: Bloomsbury Publishing.
- Ambrose, L., & Butynski, T.M. (2013a). *Galagoides demidovii* - Demidoff's dwarf galago. In T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 459-461). London: Bloomsbury Publishing.
- Ambrose, L., & Butynski, T.M. (2013b) *Galagoides thomasi* - Thomas's dwarf galago. In T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 462-466). London: Bloomsbury Publishing.
- Ambrose, L., & Oates, J.F. (2013). *Euoticus pallidus* – Needle-clawed galago. In T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 444-446). London: Bloomsbury Publishing.
- Ambrose, L., & Pimley, E.R. (2013). *Sciurocheirus alleni* – Allen's squirrel galago. In T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 418-420). London: Bloomsbury Publishing.

455 Anderson, J. (2000). Sleep-related behavioural adaptations in free-ranging anthropoid primates. *Sleep*
456 *Medicine Reviews*, 4(4), 355-373.

457 Ankel-Simons, F. (2007) *Primate Anatomy: an introduction*. Academic Press: New York.

458 Baker, S.E. (2013). Accumulation behaviours and taphonomic signatures for extant Verreaux's eagle
459 nests, *Aquila verreauxii*, in Southern Africa (Unpublished master dissertation). University of the
460 Witwatersrand, Johannesburg.

461 Bearder, S.K. (2007). A comparison of calling patterns in two nocturnal primates, *Otolemur*
462 *crassicaudatus* and *Galago moholi* as a guide to predation risk. In S. Gursky (Ed.), *Primate Anti-*
463 *predator Strategies* (pp. 206-221). Springer US.

464 Bearder, S.K., & Honess, P.E. (1992). *A survey of nocturnal primates and other mammals in the Korup*
465 *National Park, Cameroon*. Unpublished report, Oxford, UK, Nocturnal Primate Research Group.

466 Bearder, S.K., & Martin, R.D. (1980). *Acacia* gum and its use by bushbabies, *Galago senegalensis*
467 (Primates Lorisidae). *International Journal of Primatology*, 1(2), 103-128.

468 Bearder, S.K., Nekaris, K.A I., & Buzzell, C.A. (2002). Dangers in the night: are some nocturnal primates
469 afraid of the dark? In L. Miller (ed.), *Eat or be eaten: Predator sensitive foraging among primates* (pp.
470 21-43), Cambridge University Press: Cambridge.

471 Bearder, S.K., Ambrose, L., Harcourt, C., Honess, P., Perkin, A., Pimley, E., Pullen, S., & Svoboda, N.
472 (2003). Species-typical patterns of infant contact, sleeping site use and social cohesion among
473 nocturnal primates in Africa. *Folia Primatologica*, 74, 337-354.

474 Bersacola, E., Svensson, M.S., & Bearder, S.K. (2015). Niche partitioning and environmental factors
475 affecting abundance of strepsirrhines in Angola. *American Journal of Primatology*, 77, 1179-1192.

476 Bloch, J.I., & Boyer, D.M. (2002). Grasping primate origins. *Science*, 298(5598), 1606-1610.

477 Boesch, C., & Boesch, H. (1989). Hunting behavior of wild chimpanzees in the Tai National Park.
478 *American Journal of Physical Anthropology*, 78(4), 547-573.

479 Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C-H., Xie, D., Suchard, MA., Rambaut, A., &
 480 Drummond, A.J. (2014). BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS*
 481 *Computational Biology*, 10(4), e1003537.

482 Burnham, D., Bearder, S.K., Cheyne, S.M., Dunbar, R.I.M., & Macdonald, D.W. (2012). Predation by
 483 mammalian carnivores on nocturnal primates: is the lack of evidence support for the effectiveness of
 484 nocturnality as an antipredator strategy? *Folia Primatologica*, 83(3-6), 236-251.

485 Butynski, T.M. (1982). Blue monkey (*Cercopithecus mitis stuhlmanni*) predation on galagos. *Primates*, 23,
 486 563-566.

487 Butynski, T.M., & De Jong, Y.A. (2004). Natural history of the Somali lesser galago (*Galago gallarum*).
 488 *Journal of East African Natural History*, 93, 23-38.

489 Butynski, T.M., & De Jong, Y.A. (2007). Distribution of the potto (Primates: Lorisidae) *Perodicticus potto*
 490 (Müller, 1776) in eastern Africa, with a description of a new subspecies from Mount Kenya. *Journal*
 491 *of East African Natural History*, 96, 113-147.

492 Butynski, T.M., & De Jong, Y.A. (2013). *Galago gallarum* Somali lesser galago. In T.M. Butynski, J.
 493 Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 434-436). London:
 494 Bloomsbury Publishing.

495 Butynski, T.M., & De Jong, Y.A. (2014). Primate conservation in the rangeland agroecosystem of Laikipia
 496 County, central Kenya. *Primate Conservation*, 28, 117-128.

497 Butynski, T.M., & De Jong, Y.A. (2017). The Mount Kenya potto is a subspecies of the eastern potto
 498 *Perodicticus ibeanus*. *Primate Conservation*, 31, Online First.

499 Butynski, T.M., De Jong, Y.A., Perkin, A.W., Bearder, S.K., & Honess, P.E. (2006). Taxonomy,
 500 distribution, and conservation status of three species of dwarf galagos (*Galagoides*) in eastern
 501 Africa. *Primate Conservation*, 21, 63-79.

502 Campbell, S.S., & Tobler, I. (1984). Animal sleep: a review of sleep duration across phylogeny.
 503 *Neuroscience & Biobehavioral Reviews*, 8, 269-300.

504 Capellini, I., Barton R.A., McNamara, P., Preston, B.T., & Nunn, C.L. (2008). Phylogenetic analysis of the
 505 ecology and evolution of mammalian sleep. *Evolution*, 62(7), 1764-1776.

506 Charles-Dominique, P. (1977). *Ecology and behaviour of nocturnal primates*. New York: Columbia
 507 University Press.

508 Charles-Dominique, P., & Martin, R.D. (1972). *Behaviour and ecology of nocturnal prosimians: field*
 509 *studies in Gabon and Madagascar*. Berlin: P. Parey.

510 Congdon, K.A., & Ravosa, M.J. (2016). Get a grip: Substrate orientation and digital grasping pressures in
 511 strepsirrhines. *Folia Primatologica*, 87, 224-243.

512 Covert, H.H. (2002). The earliest fossil primates and the evolution of prosimians: introduction. In W.C.
 513 Hartwig (Ed.), *Cambridge studies in biological and evolutionary anthropology* (pp. 13-20).
 514 Cambridge: Cambridge University Press.

515 Daussmann, K.H., Glos, J., Ganzhorn, J.U., & Helmaier, G. (2004). Physiology: Hibernation in a tropical
 516 primate. *Nature*, 429, 825-826.

517 De Jong, Y.A., & Butynski, T.M. (2004a). Surveys of the Somali lesser bushbaby (*Galago gallarum*) and
 518 northern lesser bushbaby (*Galago senegalensis*) in Kenya and Ethiopia. Unpublished report,
 519 Nanyuki, Kenya, Eastern Africa Primate Diversity and Conservation Program.

520 De Jong, Y.A., & Butynski, T.M. (2004b). Life in the thornbush – the Somali bushbaby. *Swara*, 27, 22.

521 De Jong, Y.A., & Butynski, T.M. (2009). *Primate biogeography, diversity, taxonomy and conservation of*
 522 *the coastal forests of Kenya*. Unpublished report, Nanyuki, Kenya, Eastern Africa Primate Diversity
 523 and Conservation Program.

524 De Jong, Y.A., & Butynski, T.M. (2010). *Assessment of the primates, large mammals and birds of the*
 525 *Mathews Range Forest Reserve, central Kenya*. Unpublished report, Nanyuki, Kenya, Eastern Africa
 526 Primate Diversity and Conservation Program.

527 Di Bitetti, M.S., Vidal, E.M.L., Baldovino, M.C., & Benesovsky, V. (2000). Sleeping site preferences in
 528 tufted capuchin monkey (*Cebu paella nigritus*). *American Journal of Primatology*, 50, 257-274.

529 Doody, K.Z., Howell, K.M., & Fanning, E. (2001). *West Kilombero Scarp Forest Reserve – Zoological*
530 *Report*. Unpublished report, Dar es Salaam, Tanzania, University of Dar es Salaam.

531 Drummond, A.J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC*
532 *Evolutionary Biology*, 7, 214.

533 Duarte, M.H., & Young, R.J. (2011). Sleeping site selection by urban marmosets (*Callithrix penicillata*)
534 under conditions of exceptionally high predator density. *International Journal of Primatology*, 32, 329-
535 334.

536 Eberle, M., & Kappeler, P.M. (2004). Sex in the dark: determinants and consequences of mixed male
537 mating tactics in *Microcebus murinus*, a small solitary nocturnal primate. *Behavioral Ecology and*
538 *Sociobiology*, 57, 77-90.

539 Ehrlich, A., & Macbride, L. (1989). Mother-infant interactions in captive slow lorises (*Nycticebus coucang*).
540 *American Journal of Primatology*, 19, 217-228.

541 Elgar, M.A., Pagel, M.D., & Harvey, P.H. (1988). Sleep in mammals. *Animal Behaviour*, 36, 1407-1419.

542 Engelbrecht, D. (2016). Galagos as avian nest predators in South Africa. *Primates*, 57, 455-458.

543 Eppley, T.M., Watzek, J., Dausmann, K.H., Ganzhorn, J.U., & Donati, G. (2017). Huddling is more
544 important than rest site selection for thermoregulation in southern bamboo lemurs. *Animal*
545 *Behaviour*, 127, 153-161.

546 Fleagle, J.G. (2013). *Primate adaptation and evolution*. San Diego: Elsevier.

547 Fultz, A., Brent, L., Breaux, S.D., & Grand, A.P. (2013). An evaluation of nest-building behaviour by
548 sanctuary chimpanzees with access to forested habitats. *Folia Primatologica*, 84, 405-420.

549 Gebo, D.L., & Dagosto, M. (2004). Anthropoid origins: postcranial evidence from the Eocene of Asia. In
550 C.F. Ross & R.F. Kay (Eds.), *Anthropoid origins, new visions* (pp. 369-380). New York: Springer.

551 Génin, F., Yokwana, A., Kom, N., Couette, S., Dieuleveut, T., Nash, S.D., & Masters, J.C. (2016). A new
552 galago species for South Africa (Primates: Strepsirhini: Galagidae). *African Zoology*, 51, 135-143.

553 Grow, N., Gursky-Doyen, S. (2010). Preliminary data on the behaviour, ecology, and morphology of
 554 pygmy tarsiers (*Tarsius pumilus*). *International Journal of Primatology*, 31, 1174-1191.

555 Gursky, S. (2003). Lunar philia in a nocturnal primate. *International Journal of Primatology*, 24, 351-367.

556 Harcourt CS & Perkin AW (2013) *Otolemur garnettii* – Small-eared greater galago. In T.M. Butynski, J.
 557 Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 413-416). London:
 558 Bloomsbury Publishing.

559 Hardus, M.E., Lameira, A.R., Zulfa, A., Atmoko, S.S.U., de Vries, H., & Wich, S.A. (2012). Behavioral,
 560 ecological, and evolutionary aspects of meat-eating by Sumatran orangutans (*Pongo abelii*).
 561 *International Journal of Primatology*, 33, 287-304.

562 Hart, J.A., Katembo, M., & Punga, K. (1996). Diet, prey selection and ecological relations of leopard and
 563 golden cat in the Ituri Forest, Zaire. *African Journal of Ecology*, 34, 364-379.

564 Hohmann, G., & Fruth, B. (2008). New records on prey capture and meat eating by bonobos at Lui
 565 Kotale, Salonga National Park, Democratic Republic of Congo. *Folia Primatologica*, 79, 103-110.

566 Honess, P.E., Perkin, A.W., & Butynski, T.M. (2013). *Galagoides zanzibaricus* – Zanzibar dwarf galago. In
 567 T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 447-449).
 568 London: Bloomsbury Publishing.

569 Huelsenbeck, J.P., Nielsen, R., & Bollback, J.P. (2003). Stochastic mapping of morphological characters.
 570 *Systematic Biology*, 52, 131 – 158.

571 Kappeler, P.M. (1998). Nests, tree holes, and the evolution of primate life histories. *American Journal of*
 572 *Primatology*, 46, 7-33.

573 Kappeler, P.M., Cuzzo, F.P., Fichtel, C., Ganzhorn, J.U., Gursky-Doyen, S., Irwin, M.T., Ichino, S.,
 574 Lawler, R., Nekaris, K.A.I., Ramanamanjato, J.B. (2017) Radespiel, U. Long-term field studies of
 575 lemurs, lorises, and tarsiers. *Journal of Mammalogy*, 98, 661-669.

576 Katoh, K., & Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7:
 577 improvements in performance and usability. *Molecular Biology and Evolution*, 30, 772-780.

578 Kenyon, M., Streicher, U., Loung, H., Tran, T., Tran, M., Vo, B., & Cronin, A. (2014). Survival of
579 reintroduced pygmy slow loris *Nycticebus pygmaeus* in south Vietnam. *Endangered Species*
580 *Research*, 25, 185-195.

581 Kingdon, J. (2015). *The Kingdon field guide to African mammals*. London: Bloomsbury.

582 Kumara, H.N., Sasi, R., Chandran, S. & Radhakrishna, S. (2016). Distribution of the grey slender loris
583 (*Loris lydekkerianus* Cabrera, 1908) in Tamil Nadu, southern India. *Folia Primatologica*, 87, 291–
584 302.

585 Lesku, J. A., Roth II, T.C., Amlaner, C.J., & Lima, S.L. (2006). A phylogenetic analysis of sleep
586 architecture in mammals: the integration of anatomy, physiology, and ecology. *The American*
587 *Naturalist*, 168, 441-453.

588 Lima, S.L., Rattenborg, N.C., Lesku, J.A. & Amlaner, C.J. (2005). Sleeping under the risk of predation.
589 *Animal Behavior*, 70, 723-736.

590 Llorente, M., Sabater, Pi. J., & Houle, A. (2003). Association between *Galago thomasi* and *Pan*
591 *troglodytes schweinfurthii* in the Kibale National Park, Uganda. *Folia Primatologica*, 74, 80-84.

592 Lock, L.C., & Anderson, J.R. (2013). Kin, daytime associations, or preferred sleeping sites? Factors
593 influencing sleep site selection in captive chimpanzees (*Pan troglodytes*). *Folia Primatologica*, 84,
594 158-169.

595 Lumsden, W.H.R., & Masters, J. (2001). Galago (Galagonidae) collections in East Africa (1953-1955):
596 ecology of the study areas. *African Primates*, 5, 37-42.

597 Lutermann, H., Verburgt, L., & Rendigs, A. (2010). Resting and nesting in a small mammal: sleeping sites
598 as a limiting resource for female grey mouse lemurs. *Animal Behaviour*, 79, 1211-1219.

599 Masters, J.C., Génin, F., Couette, S., Groves, C.P., Nash, S.D., Delpero, M., & Pozzi, L. (2017). A new
600 genus for the eastern dwarf galagos (Primates: Galagidae). *Zoological Journal of the Linnaean*
601 *Society*, 181, 229-241.

602 McGrew, W.C., Tutin, C.E.G., & Baldwin, P.J. (1978). Primates preying upon vertebrates: new records
603 from West Africa (*Pan troglodytes verus*, *Papio papio*, *Cercopithecus sabaeus*). *Carnivore*, 1, 41-45.

604 Mueller, A.E., & Thalmann, U. (2000). Origin and evolution of primate social organization: a
 605 reconstruction. *Biological Reviews of the Cambridge Philosophical Society*, 75, 405-435.

606 Munds, R.A., Nekaris, K.A.I., & Ford, S.M. (2013). Taxonomy of the Bornean slow loris, with new species
 607 *Nycticebus kayan* (Primates, Lorisidae). *American Journal of Primatology*, 75, 46-56.

608 Msuya, C.A. (1993). Feeding habits of crowned eagles (*Stephanoaetus coronatus*) in Kiwengoma Forest
 609 Reserve, Matumbi Hills, Tanzania. *Annales Musee Royal de l'Afrique Centrale. Sciences*
 610 *Zoologiques*, 118-120.

611 Mzilikazi, N., Masters, J.C., & Lovegrove, B.G. (2006). Lack of torpor in free-ranging southern lesser
 612 galagos, *Galago moholi*: ecological and physiological considerations. *Folia Primatologica*, 77, 465-
 613 476.

614 Nash, L., Zimmermann, E., & Butynski, T.M. (2013). *Galago senegalensis* - Northern lesser galago. In
 615 T.M. Butynski, J. Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 425-429).
 616 London: Bloomsbury Publishing.

617 Nekaris, K.A.I. (2003a). Spacing system of the Mysore slender loris (*Loris lydekkerianus*
 618 *lydekkerianus*). *American Journal of Physical Anthropology*, 121, 86-96.

619 Nekaris, K.A.I. (2003b). Observations on mating, birthing and paternal care in three taxa of slender loris in
 620 India and Sri Lanka (*Loris tardigradus* and *Loris lydekkerianus*). *Folia Primatologica*, 74, 312-336.

621 Nekaris, K.A.I. (2013a). Galagidae. In R.A. Mittermeier, A.B. Rylands, & D.E. Wilson (Eds.), *Handbook of*
 622 *the Mammals of the World: 3. Primates* (pp. 184-209). Barcelona: Lynx Ediciones.

623 Nekaris, K.A.I. (2013b). Lorisidae. In R.A. Mittermeier, A.B. Rylands, & D.E. Wilson (Eds.), *Handbook of*
 624 *the Mammals of the World: 3. Primates* (pp. 210-235). Barcelona: Lynx Ediciones.

625 Nekaris, K.A.I. (2014). Extreme primates: ecology and evolution of Asian lorises. *Evolutionary*
 626 *Anthropology* 23, 177-187.

627 Nekaris, K.A.I., & Bearder, S.K. (2011). The strepsirrhine primates of Asia and mainland Africa: diversity
 628 shrouded in darkness. In C. Campbell, A. Fuentes, K. MacKinnon, S.K. Bearder, & R. Stumpf (Eds.),
 629 *Primates in Perspective 2nd edition* (pp. 34-54). Oxford: Oxford University Press.

630 Nekaris, K.A.I., & Jayewardene, J. (2003). Pilot study and conservation status of the slender loris (*Loris*
 631 *tardigradus* and *L. lydekkerianus*) in Sri Lanka. *Primate Conservation*, 19, 83-90.

632 Nekaris, K.A.I., Poindexter, S.A., Reinhardt, K.D., Sigaud, M., Cabana, F., Wirdateti, W., & Nijman, V.
633 (2017). Coexistence between Javan slow lorises (*Nycticebus javanicus*) and humans in a dynamic
634 agroforestry landscape in West Java, Indonesia. *International Journal of Primatology*, 38, 303-320.

635 Nishida, T., Uehara, S., & Nyundo, R. (1979). Predatory behavior among wild chimpanzees of the Mahale
636 Mountains. *Primates*, 20, 1-20.

637 Nowack, J., Mzilikazi, N., & Dausmann, K.H. (2010). Torpor on demand: heterothermy in the non-lemur
638 primate *Galago moholi*. *PLoS One*, 5(5), p.e10797.

639 Nowack, J., Mzilikazi, N., & Dausmann, K.H. (2013a). Torpor as an emergency solution in *Galago moholi*:
640 heterothermy is triggered by different constraints. *Journal of Comparative Physiology B*, 183, 547-
641 556.

642 Nowack, J., Wippich, M., Mzilikazi, N., & Dausmann, K.H. (2013b). Surviving the cold, dry period in Africa:
643 behavioral adjustments as an alternative to heterothermy in the African lesser bushbaby (*Galago*
644 *moholi*). *International Journal of Primatology*, 34, 49-64.

645 Oates, J.F. (1984). The niche of the Potto, *Perodicticus potto*. *International Journal of Primatology*, 5, 51-
646 61.

647 Off, E.C., Isbell, L.A., & Young, T.P. (2008). Population density and habitat preferences of the Kenya
648 lesser galago (*Galago senegalensis braccatus*) along the Ewaso Nyiro River, Laikipia, Kenya.
649 *Journal of East African Natural History*, 97, 109-116.

650 O'Malley, R.C. (2010). Two observations of galago predation by the Kasakela Chimpanzees of Gombe
651 Stream National Park, Tanzania. *Pan-Africa News*, 17, 17-19.

652 Peckre, L., Fabre, A.C., Wall, C.E., Brewer, D., Ehmke, E., Haring, D., Shaw, E., Welser, K., &
653 Pouydebat, E. (2016). Holding-on: co-evolution between infant carrying and grasping behaviour in
654 strepsirrhines. *Scientific Reports*, 6: 37729.

655 Perkin, A.W. (2000). *The taxonomic status of bushbabies (galagos) in the Uluguru Mountains*.
656 Unpublished report, Dar es Salaam, Tanzania, Tanzania Forest Conservation Group.

657 Pimley, E.R. (2002). *The behavioural ecology and genetics of two nocturnal prosimians: pottos*
658 (*Perodicticus potto edwardsi*) and *Allen's bushbabies* (*Galago alleni cameronsensis*) (Unpublished
659 doctoral dissertation). University of Cambridge, Cambridge.

660 Pimley, E.R., & Bearder, S.K. (2013). *Perodicticus potto* – Potto. In T.M. Butynski, J. Kingdon & J. Kalina
 661 (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 393-399). London: Bloomsbury Publishing.

662 Pimley, E.R., Bearder, S.K., & Dixon, A.F. (2005a). Social organization of the Milne-Edward's potto.
 663 *American Journal of Primatology*, 66, 317-330.

664 Pimley, E.R., Bearder, S.K., & Dixon, A.F. (2005b). Home range analysis of *Perodicticus potto* and
 665 *Sciurocheirus cameronensis*. *International Journal of Primatology*, 26, 191-205.

666 Pontes, A.R.M., & Soares, M.L. (2005). Sleeping sites of common marmosets (*Callithrix jacchus*) in
 667 defaunated urban forest fragments: a strategy to maximize food intake. *Journal of Zoology*, 266, 55-
 668 63.

669 Pozzi, L., Nekaris, K.A.I., Perkins, A., Bearder, S.K., Pimley, E.R., Schulze, H., Streicher, U., Nadler, T.,
 670 Kitchener, A., Zischler, H., Zinner, D., & Roos, C. (2015). Remarkable ancient divergences amongst
 671 neglected lorisiform primates. *Zoological Journal of the Linnaean Society*, 175, 661-674.

672 Pruetz, J.D., & Bertolani, P. (2007). Savanna chimpanzees, *Pan troglodytes verus*, hunt with tools.
 673 *Current Biology*, 17, 412-417.

674 Pruetz, J.D., Bertolani, P., Ontl, K.B., Lindshield, S., Shelley, M., & Wessling, E.G. (2015). New evidence
 675 on the tool-assisted hunting exhibited by chimpanzees (*Pan troglodytes verus*) in a savannah habitat
 676 at Fongoli, Sénégal. *Royal Society Open Science*, 2, 140507.

677 Pullen, S., & Bearder, S.K. (2013). *Galago moholi* – Southern lesser galago. In T.M. Butynski, J.
 678 Kingdon & J. Kalina (Eds.), *Mammals of Africa* (Volume II: Primates, pp. 430-433). London:
 679 Bloomsbury Publishing.

680 Qihai, Z., Chengming, H., Ming, L., & Fuwen, W. (2009). Sleeping site use by *Trachypithecus*
 681 *francoisi* at Nonggang Nature Reserve, China. *International Journal of Primatology*, 30, 353-
 682 365.

683 Radespiel, U., Cepok, S., Zietemann, V. & Zimmermann, E. (1998). Sex-specific usage patterns of
 684 sleeping-sites in grey mouse lemurs. *American Journal of Primatology*, 46, 77-84.

685 Rasmussen, D.T. (1986). *Life history and behavior of slow lorises and slender lorises: implications for the*
 686 *lorisine-galagine divergence* (Unpublished doctoral dissertation). Duke University, USA.

687 Rattenborg, N.C., Martinez-Gonzalez, D., & Lesku, J.A. (2009). Avian sleep homeostasis: convergent
688 evolution of complex brains, cognition and sleep functions in mammals and birds. *Neuroscience &*
689 *Biobehavioral Reviews*, 33, 253-270.

690 Ray, I., Wren, B.T., & Bowers, E.J. (2016). Documentation of plant consumption by *Galago moholi* in
691 South Africa. *African Primates*, 11, 45-48.

692 Reinhardt, K.D., Wirdateti, & Nekarlis, K.A.I. (2016) Climate-mediated activity of the Javan slow loris,
693 *Nycticebus javanicus*. *AIMS Environmental Science*, 3, 249-260.

694 Revell, L.J. (2012) Phytools: an R package for phylogenetic comparative biology (and other things).
695 *Methods in Ecology and Evolution*, 3, 217 – 223.

696 Rovero, F., Marshall, A.R., Jones, T., & Perkin, A. (2009). The primates of the Udzungwa Mountains:
697 diversity, ecology and conservation. *Journal of Anthropological Sciences*, 87, 93-126.

698 Ruf, T., Streicher, U., Stalder, G.L., Nadler, T., & Walzer, C. (2015). Hibernation in the pygmy slow loris
699 (*Nycticebus pygmaeus*): multiday torpor in primates is not restricted to Madagascar. *Scientific*
700 *Reports*, 5, 17392.

701 Sabater, Pi.J., Veá, J.J., & Serrallonga, J. (1997). Did the first hominids build nests? *Current*
702 *Anthropology*, 38, 914-916.

703 Samson, D.R., & Nunn, C.L. (2015). Sleep intensity and the evolution of human cognition. *Evolutionary*
704 *Anthropology* 24, 225-237.

705 Samson D.R., & Shumaker R.W. (2013). Documenting orang-utan sleep architecture: sleeping platform
706 complexity increases sleep quality in captive *Pongo*. *Behaviour*, 150, 845-861.

707 Samson D.R., & Shumaker R.W. (2015a). Orangutans (*Pongo* spp.) have deeper, more efficient sleep
708 than baboons (*Papio papio*) in captivity. *American Journal of Primatology*, 157, 421-427.

709 Samson, D.R., & Shumaker, R.W. (2015b). Pre-sleep and sleeping platform construction behaviour in
710 captive orangutans (*Pongo* spp.): Implications for ape health and welfare. *Folia Primatologica*, 86,
711 187-202.

712 Schmid, J. (1998). Tree holes used for resting by gray mouse lemurs (*Microcebus murinus*) in
713 Madagascar: Insulation capacities and energetic consequences. *International Journal of*
714 *Primatology*, 19, 797-809.

715 Schülke, O., & Ostner, J. (2001). Predation on *Lepilemur* by a harrier hawk and implications for sleeping
716 site quality. *Lemur News*, 6, 5.

717 Sellers W.I. (1996). A biomechanical investigation into the absence of leaping in the locomotor repertoire
718 of the slender loris (*Loris tardigradus*). *Folia Primatologica*, 67, 1-14.

719 Shultz, S., Noë, R., McGraw, W.S., & Dunbar, R.I.M. (2004). A community-level evaluation of the impact
720 of prey behavioural and ecological characteristics on predator diet composition. *Proceedings of the*
721 *Royal Society B: Biological Sciences*, 271, 725-732.

722 Suchard, M.A. & Rambaut, A. (2009). Many-Core Algorithms for Statistical Phylogenetics. *Bioinformatics*,
723 25, 1370-1376.

724 Svensson, M.S., & Bearder, S.K. (2013). Sightings and habitat use of the northern lesser galago (*Galago*
725 *senegalensis senegalensis*) in Niimi National Park, The Gambia. *African Primates*, 8, 51-58.

726 Svensson, M.S., Bersacola, E., Mills, M.S.L., Munds, R.A., Nijman, V., Perkin, A., Masters, J.C., Couette,
727 S., Nekaris, K.A.I., & Bearder, S.K. (2017). A giant among dwarfs: a new species of galago
728 (Primates: Galagidae) from Angola. *American Journal of Physical Anthropology*, 163, 30-43.

729 Tagg, N., Willie, J., Petre, C.A., & Haggis, O. (2013). Ground night nesting in chimpanzees: New insights
730 from Central chimpanzees (*Pan troglodytes troglodytes*) in south-east Cameroon. *Folia*
731 *Primatologica*, 84, 362-383.

732 Wiens, F., & Zitzmann, A. (1999). Predation on a wild slow loris (*Nycticebus coucang*) by a reticulated
733 python (*Python reticulatus*). *Folia Primatologica*, 70, 362-364.

734 Uehara, S. (1997). Predation on mammals by the chimpanzee (*Pan troglodytes*). *Primates*, 38, 193-214.

735 Utami, S.C., & van Hooff, J.A.R.A.M. (1997). Meat-eating by adult female Sumatran orangutans (*Pongo*
736 *pygmæus abelii*). *American Journal of Primatology*, 43, 159-165.

737 Voskamp A., Rode E.J., Coudrat C.N., Wilson R.J., & Nekaris K.A.I. (2014). Modelling the habitat use and
738 distribution of the threatened Javan slow loris *Nycticebus javanicus*. *Endangered Species Research*,
739 23, 277-286.

740 Zepelin, H. (2000). Mammalian sleep. In M.H. Kryger, T. Roth & W.C. Dement (Eds.), *Principles and*
741 *Practice of Sleep Medicine*. (pp. 82-92). Philadelphia: W. B. Saunders.

FIGURE 1 Ancestral state reconstructions of stochastic character mapping of lorisiform fur-clinging whereby infants cling onto the fur of their parents when being transported. Red numbers indicate the Bayesian posterior probabilities of the phylogenetic tree if <1. Branches above nodes (closer to tips) are coloured based on their ancestral state probability. Pie charts on nodes and black numbers (states 1/2/3) indicate the probability of the state in the common ancestor. The states are in the following order: 1 = absent, 2 = occasionally present, 3 = present.

FIGURE 2 Ancestral state reconstructions of stochastic character mapping of lorisiform sleep site use: a) bamboo, b) branch, c) dense tangle, d) nest, e) tree hole. Branches above nodes (closer to tips) are coloured based on their ancestral state probability. Pie charts on nodes and black numbers (states 1/2/3) indicate the probability of the state in the common ancestor. The states are in the following order: 1 = absent, 2 = occasionally present, 3 = present; except for the branch sleep site where: 1 = absent, 2 = present.

FIGURE 3 Young chimpanzee *Pan troglodytes* in Guinea holding a dead northern lesser galago *Galago senegalensis*, having caught it in the daytime. Photo by: Chimpanzee Conservation Center / Charlotte Houpline.

TABLE 1 Sleep site type: 0 - no evidence of use, 1 -irregular or occasional use or mixed evidence use from different studies, 2 - regular or habitual use of nests, ? - evidence is based on anecdotal information or when information is lacking, * - using man-made structures as sleep sites Social cohesion: 0 – no, 1 – yes.

TABLE 2 Day-time predation of lorisiforms by great apes







769

770

TABLE 1 Sleep site type: 0 - no evidence of use, 1 -irregular or occasional use or mixed evidence use from different studies, 2 - regular or habitual use of nests, ? - evidence is based on anecdotal information or when information is lacking, * - using man-made structures as sleep sites Social cohesion: 0 – no, 1 – yes.

Species				Sleep site type					Social cohesion				Non-human diurnal and crepuscular predators (Confirmed records in bold)	Predator avoidance strategies	Reference
	Intermembral Index	Mouth-carrying	Fur-clinging	Nest	Tree hole	Dense tangle	Branch/fork	Bamboo	Audible advertising calls	Audible contact maintenance calls	Allogrooming at sleep site	Sleep group size (range)			
<i>Galagoides demidovii</i>	68	2	0	2	1	1	0	1	1	1	?	2-10	Sooty mangabeys, bonobos		1; 2; 3; 4 ; 5; 6
<i>Gd. thomasi</i>	67	2	0	1	1	1	1	0	1	1	?	<5	Snakes, hawks, hornbill, viverrids, mongooses, blue monkey		7; 8; 9
<i>Gd. kumbirensis</i>		?	?	?	?	?	?	?	?	?	?	?	?		10
<i>Paragalago orinus</i>		1	0	2	1	1	?	0	1	?	1-5	1-9	Snakes, genets, Sykes's monkeys		11;12; 13
<i>Pg. rondoensis</i>		2	0	2	0	?	?	?	1	0	?	<3	Snakes e.g. forest cobras, boomslangs, green mambas)		13
<i>Pg. granti</i>		2	0	2	2	0	0	0	1	0	?	4-5	Snakes e.g. forest cobras, boomslangs, green mambas)		13; 14; 15
<i>Pg. cocos</i>		2	0	0	2*	0	0	0	1	1	?	1-5	Snakes e.g. forest cobras, boomslangs, green mambas), <i>Cercopithecus</i> monkeys		16; 17
<i>Pg. zanzibaricus</i>	60	2	0	2	2	2	0	0	1	1	?	1-5	Snakes e.g. forest cobras, boomslangs, green mambas)		13; 16; 17; 18
<i>Galago senegalensis</i>	52	2	0	1	2*	2	1	0	1	1	1	<6	Chimpanzees		19; 20; 21; 22; 23; 24; 25; 26; 27; 28; 29;30
<i>G. gallarum</i>		?	0	1	1	2	2	0	1	?	1	≤3	Snakes, raptors, jackals, mongooses, genets, wild cats	Live in the thorniest habitats of all galagos. Sleep and rest in the core of thorny vegetation.	17; 31; 32; 33; 34; 35
<i>G. moholi</i>	54	2	0	2	2*	1	1	0	1	0	?	1-8.	Large snakes & monitor lizards, genets, Verreaux's eagle , small carnivores,		13; 36; 37; 38; 39; 40; 41
<i>G. matschiei</i>		2	0	?	2	?	?	?	1	0	?	?	Large snakes, viverrids, blue monkeys , baboons, chimpanzees		7; 42
<i>Sciurocheirus alleni</i>	65	2	0	1	2	2	1	?	1	1	1	1-6	Large snake, viverrids		1; 16; 43; 44
<i>S. gabonensis</i>		2	0	0	2	?	0	?	1	?	?	1, 1-3	Large snakes, viverrids, leopard, African golden cat		16; 45
<i>S. cameronensis</i>		2	0	1	2	?	?	?	?	?	?	1, 1-6			16; 44; 46
<i>S. makandensis</i>		?	0	?	?	?	?	?	1	1	?	1-4	Large snakes, viverrids, golden cats		47; 48

<i>Euoticus elegantulus</i>	64	0	2	0	0	1?	1?	0	1	1	?	3-4	Pythons, viverrids	1; 13; 16	
<i>E. pallidus</i>		0	2	0	2	0	1	0	1	1	?	1-4	Central African linsang	13; 16; 49	
<i>Otolemur crassicaudatus</i>	70	1	1	1	1*	2	0	?	1	0	?	1-4	Large snakes, raptors, leopards, chimpanzees	13; 16; 17; 50	
<i>O. garnettii</i>	69	1	1	0	1	2	?	?	1	1	?	1-4	Large snakes, raptors	17; 51; 52; 53	
<i>Arctocebus calabarensis</i>	89	0	2	0	0	2	1	0	0	0	?	1-2	Snakes, viverrids, monkeys	13	
<i>A. aureus</i>		0	2	0	0	2	1	0	0	0	?	1-2	Large snakes, viverrids,	13	
<i>Perodicticus potto</i>	88	0	2	0	0	2	1	0	0	0	?	?	Large snakes, African crowned eagles , civets, black-legged mongoose, leopards , African golden cats, <i>Cercopithecus</i> monkeys, mandrills, chimpanzees	Scapular neck shield, predator defense posture, drops to ground	6; 19; 54; 55; 56; 57
<i>P. edwardsi</i>		0	2	0	?	2	?	0	0	0	?	1-2		Scapular neck shield, predator defense posture, drops to ground	43; 58; 59
<i>P. ibeanus</i>		0	2	?	?	2	?	0	0	0	?	?	Leopard	Scapular neck shield, predator defense posture, drops to ground	54; 5
<i>Nycticebus javanicus</i>	93	0	2	0	0	2	1	2	0	0	1	4		Venomous, predator defense posture	60
<i>N. bengalensis</i>		0	2	0	0	2	1	2	0	0	1	?		Venomous, sleeps high in trees inaccessible positions or in dense thorny tangles	61
<i>N. menagensis</i>	91	0	2	0	0	2	1	0	0	0	1	1-3	Reticulated pythons , raptors	Venomous, predator defense posture	62
<i>N. pygmaeus</i>	91	0	2	0	0	2	1	1	1	1	1	2-5	Monitor lizards, raptor, small carnivores	Venomous, predator defense posture	63; 64
<i>N. coucang</i>	91	0	2	0	0	2	1	1	0	1	1	3	Reticulated python, monitor lizards	Venomous, predator defense posture	65; 66
<i>N. hilleri</i>	89	0	2	0	0	2	1	1	1	1	?	?	Changeable hawk eagle, Sumatran orangutans	Venomous, predator defense posture	67; 68; 69; 70
<i>N. kayan</i>		?	?	?	?	?	?	?	?	?	?	?		Venomous, predator defense posture	
<i>N. bancanus</i>		?	?	?	?	?	?	?	?	?	?	?	?	?	
<i>N. borneanus</i>		?	?	?	?	?	?	?	?	?	?	?	?	Venomous, predator defense posture	
<i>Loris tardigradus</i>	90	0	2	0	0	2	1	0	1	1	1	4		Sleeps in dense tangles, has cobra defense posture	71
<i>L. lydekkerianus</i>	92	0	2	0	0	2	1	0	1	1	1	4-5	Rusty spotted cat	Sleeps in dense tangles, has cobra	72; 73

1) Charles Dominique, 1977; **2)** Bearder & Honess, 1992; **3)** Hohmann & Fruth, 2008; **4)** Ambrose & Butynski 2013a; **5)** A. Luhrs, pers. obs.; **6)** E. Pimley, pers. obs.; **7)** Butynski, 1982; **8)** Llorente et al., 2003; **9)** Ambrose & Butynski 2013b; **10)** Svensson et al., 2017; **11)** Perkin, 2000; **12)** Doody et al., 2001; **13)** Nekaris & Bearder, 2011; **14)** Butynski et al., 2006; **15)** Génin et al., 2016; **16)** Kingdon, 2015; **17)** Y. De Jong & T. Butynski, pers. obs.; **18)** Honess, Perkin & Butynski, 2013; **19)** McGrew, Tutin & Baldwin, 1978; **20)** Nishida et al., 1979; **21)** Uehara, 1997; **22)** Pruetz & Bertolani, 2007; **23)** Off, Isbell & Young, 2008; **24)** De Jong & Butynski 2009; **25)** O'Malley, 2010; **26)** Nash, Zimmermann & Butynski, 2013; **27)** Svensson & Bearder, 2013; **28)** Butynski & De Jong, 2014; **29)** Butynski & De Jong, 2017; **30)** Pruetz et al., 2015; **31)** Butynski & De Jong 2004; **32)** Butynski & De Jong, 2013; **33)** De Jong & Butynski ,2004a; **34)** De Jong & Butynski, 2004b; **35)** De Jong & Butynski, 2010; **36)** Mzilikazi, Masters & Lovegrove, 2006; **37)** Nowack, Mzilikazi & Dausmann, 2010; **38)** Burnham et al., 2012; **39)** Baker, 2013; **40)** Nowack, Wippich, Mzilikazi, & Dausmann, 2013; **41)** Pullen & Bearder, 2013; **42)** Ambrose, 2006; **43)** Pimley, 2002; **44)** Ambrose & Pimley 2013; **45)** Ambrose, 2013; **46)** Nekaris, 2013b; **47)** Ambrose, 2003; **48)** Ambrose, 2013; **49)** Ambrose & Oates, 2013; **50)** Rovero et al., 2009; **51)** Lumsden & Masters, 2001; **52)** De Jong & Butynski, 2009; **53)** Harcourt & Perkin, 2013; **54)** Hart, Katembo & Punga, 1996; **55)** Msuya, 2003; **56)** Shultz et al., 2004; **57)** Pimley & Bearder, 2013; **58)** Pimley et. al., 2005a; **59)** Pimley et. al., 2005b; **60)** Nekaris et al., 2017; **61)** N. Das, K.A.I. Nekaris & S.A. Poindexter, pers. obs.; **62)** D.J. Stark, pers. obs.; **63)** Kenyon et al., 2014; **64)** K.A.I. Nekaris, S.A. Poindexter & D.J. Stark, pers. obs.; **65)** Wiens & Zitzmann, 1999; **66)** R. Moore, pers. comm.; **67)** Utami & van Hooft, 1997; **68)** Hardus et al., 2012; **69)** C. Schuppli, pers. comm.; **70)** K.A.I. Nekaris & V. Nijman, pers. obs.; **71)** Nekaris & Jayewardene, 2003; **72)** Bearder et al., 2002; **73)** Nekaris, 2003b

TABLE 2 Day-time predation of loriforms by great apes

Species	Site, country	Predation	Reference
<i>Pan paniscus</i>	Lui Kotale, DR Congo	<i>Galagoide demidovii</i> forced out of tree hole	Hohmann & Fruth, 2008
<i>P. troglodytes</i>	Gombe Stream, Tanzania	<i>Galago senegalensis</i> retrieved from trunk and consumed	O'Malley, 2010
<i>P. troglodytes</i>	Fongoli, Senegal	<i>Galago senegalensis</i> , frequently hunted, chimpanzees using tools	Pruetz & Bertolani, 2007; Pruetz et al., 2015
<i>P. troglodytes</i>	Mt. Assirik, Senegal	<i>Galago senegalensis</i> and <i>Perodicticus potto</i> , remains found in fecal samples	McGrew et al., 1978
<i>P. troglodytes</i>	Mahale Mountains, Tanzania	<i>Otolemur crassicaudatus</i> and <i>Galago spp.</i> harassed and consumed by chimpanzees	Nishida et al., 1979; Uehara, 1997
<i>P. troglodytes</i>	Haute Niger, Guinea	Galagos . Chimpanzees observed killing, but not consuming, galagos.	C. Colin, pers. comm.
<i>P. troglodytes</i>	Ngogo / Kanyawara, Kibale, Uganda	<i>Galago spp.</i> and <i>Perodicticus ibeanus</i> . Interactions with galagos frequently observed.	J. Negrey, pers. comm.; R. Wrangham, pers. comm.
<i>P. troglodytes</i>	Bossou forest, Guinea	<i>Perodicticus potto</i> harassed by chimpanzees	K. Hockings, pers. comm.
<i>P. troglodytes</i>	Tai, Ivory Coast	<i>Perodicticus potto</i> , females and offspring observed to hunt and feed on pottos	L. Luncz, pers. obs.; E. Pimley, pers. obs.
<i>Pongo abelii</i>	Ketambe, Indonesia	<i>Nycticebus hilleri</i> , observations of orangutans eating slow lorises	Utami & van Hooff, 1997; Hardus et al., 2012; S.S. Utami Atmoko, pers. comm., S. Rimba, pers. Comm.
<i>P. abelii</i>	Suaq, Indonesia	<i>Nycticebus hilleri</i> , observations of orangutans harassing and eating slow lorises	C. Schuppli, pers. comm.